

A GEOMETRIC CHARACTERIZATION OF EXTREMAL SETS IN ℓ_p SPACES

by Viet NguyenKhac ¹ & Khiem NguyenVan ²

Hanoi Institute of Mathematics & Hanoi University of Education

1991 Mathematics Subject Classification. Primary 46B20, 46E30.

Abstract

We give a geometric characterization of extremal sets in ℓ_p spaces that generalizes our previous result for such sets in Hilbert spaces.

1. Introduction

Let $(X, \|\cdot\|)$ be a Banach space. For a non-empty bounded subset A of X and a non-empty subset B of X let's fix the following notations: $d(A) := \sup \{\|x - y\| : x, y \in A\}$ – the diameter of A ; $r_B(A) := \inf_{y \in B} \sup_{x \in A} \|x - y\|$ – the relative Chebyshev radius of A with respect to B ; in particular $r(A) := r_{\overline{\text{co}}A}(A)$ with $\overline{\text{co}}A$ denoting the closed convex hull of A ; a point $y \in B$ is called a Chebyshev center of A in B , if $\sup_{x \in A} \|x - y\| = r_B(A)$.

The Jung constant of X is defined by $J(X) := \sup \{r_X(A) : A \subset X, \text{ with } d(A) = 1\}$. The problem of estimating Jung's constant plays an important role in the geometry of Banach spaces (*cf.* [1], [7]). It is well-known that for inner-product spaces $J(E^n) = \sqrt{\frac{n}{2(n+1)}}$ ([5], *cf.* [2], [4]) and $J(H) = \frac{1}{\sqrt{2}}$ (H denotes a Hilbert space) ([8]). In general if X is an n -dimensional normed space, then $J(X) \leq \frac{n}{n+1}$. Furthermore the equality is attained for certain spaces (see [7]). As for ℓ_p , L_p spaces ($p > 1$) S. A. Pichugov ([7]) has obtained the

¹Institute of Mathematics, P.O.Box 631 Bo Ho, 10000 Hanoi, Vietnam; e-mail: nkviet@thevinh.ncst.ac.vn

²Department of Mathematics and Informatics, Hanoi University of Education, Cau Giay dist., Hanoi, Vietnam.

exact values for the Jung constant of these spaces: $J(\ell_p) = J(L_p) = \frac{1}{\sqrt[p]{2}}$, $q := \frac{p}{p-1}$, if $1 < p \leq 2$, and $J(\ell_p) = J(L_p) = \frac{1}{\sqrt[p]{2}}$, if $p > 2$.

Definition 1.1 ([6]). We say that a bounded subset A of X consisting of at least two points is an extremal set, if $r_X(A) = J(X).d(A)$.

The main result of [6] states that a bounded subset A of a Hilbert space H with $r(A) = 1$ is extremal if and only if for every $\varepsilon \in (0, \sqrt{2})$, for every positive integer m there exists an m -simplex Δ with its vertices in A and each edge of Δ has length not less than $\sqrt{2} - \varepsilon$. Furthermore for such a subset A we have $\alpha(A) = \sqrt{2}$ and $\chi(A) = 1$, where $\alpha(A)$, $\chi(A)$ denote the Kuratowski and Hausdorff measures of non-compactness of A respectively.

Our aim in this paper is to treat the next interesting case: the case of ℓ_p spaces ($1 < p < \infty$). We obtain a partial generalization of the result above. More precisely, if A is an extremal subset of a given ℓ_p space ($1 < p < \infty$), then $\alpha(A) = d(A)$. As an immediate consequence one obtains a Gulevich-type result for ℓ_p spaces: extremal sets in ℓ_p , ($1 < p < \infty$) are not relatively compact (cf. [3]). Moreover for every $\varepsilon \in (0, d(A))$, for every positive integer m there exists an m -simplex Δ with its vertices in A and each edge of Δ has length not less than $d(A) - \varepsilon$. The proof is based on a further development of a purely combinatorial method in our previous paper [6] which essentially relies on a very deep part of convex analysis. It should be noted that this observation was first noted in [4], §10.2, where the authors exposed classical Jung's theorem from the point of view of "subdifferentials", and was later extended in [7] to the case of ℓ_p spaces.

The paper is organized as follows. In §2 we review some facts related to the "Clearance"-type Theorem ("Decomposition Theorem" in translation), especially for ℓ_p spaces, we shall need in the sequel. The heart of this section is Proposition 2.1 which should be considered as an infinite-dimensional variation of the main claim in [7]. In §3 we first formulate without proof two auxiliary inequalities (Lemma 3.1) one of which was essentially due to N. I. Chernykh (personal communication to the author of [7]). Our main results are Theorems 3.2 and 3.4 the proof of which is proceeded then by means of the combinatorial method of [6] extended to ℓ_p spaces together with Proposition 2.1 and inequalities from Lemma 3.1.

2. Preliminaries

For the basic definitions and concepts in convex analysis we refer the reader to [4]. The following proposition is a slight generalization of the main proposition of [7].

Proposition 2.1. *Let X be a uniformly smooth Banach space, $A = \{x_1, x_2, \dots, x_n\}$ a finite subset with $r := r_X(A) > 0$, c a Chebyshev center of A in X . Then there exist points*

$y_1, y_2, \dots, y_m$ ($m \leq n$) in A , linear functionals $f_1, f_2, \dots, f_m$ in X^* and positive numbers $\alpha_1, \alpha_2, \dots, \alpha_m$ satisfying:

- (i) $\langle y_i - c, f_i \rangle := f_i(y_i - c) = \|y_i - c\| = r$ for $i = 1, 2, \dots, m$;
- (ii) $\|f_i\|_{X^*} = 1, \quad i = 1, 2, \dots, m$;
- (iii) $\sum_{i=1}^m \alpha_i f_i = 0, \quad \sum_{i=1}^m \alpha_i = 1$.

Proof. For completeness we give here a proof valid in all dimensions. From the uniform smoothness of X it follows that X is reflexive (hence there exists a Chebyshev center of A in X) and the mapping

$$J: X \longrightarrow 2^{X^*}$$

$$x \longmapsto J(x) := \{x^* \in X^* : \|x^*\| = 1, \langle x, x^* \rangle = \|x\|\}$$

is single-valued. Let's consider the following two functionals:

$$F: X \times A \longrightarrow \mathbb{R}$$

$$(x, a) \longmapsto F(x, a) := \|x - a\|,$$

and

$$f: X \longrightarrow \mathbb{R}$$

$$x \longmapsto f(x) := \max_{a \in A} F(x, a).$$

With $A_0(x) := \{a \in A : F(x, a) = f(x)\}$ we see that the hypotheses of Theorem 3, §4.2, Chapter 4 in [4] are fulfilled. Hence for every $x \in X$ we have

$$\overline{co} \left(\bigcup_{a \in A_0(x)} \partial F(x, a) \right) = \partial f(x) \quad (2.1)$$

where $\partial F(x, \cdot)$ and $\partial f(x)$ denote subdifferentials of F and f at x , and the closure in (2.1) is taken in the w^* -topology of the space X^* , which clearly coincides with the w -topology of X^* , since X is reflexive.

Recall that for $c \in X$ to be a Chebyshev center of A in X it is necessary and sufficient that $0 \in \partial f(c)$ (cf. [4], §1.3, Proposition 1). One may write $A_0(c) = \{y_1, y_2, \dots, y_m\}$ with $\|y_i - c\| = r$ for $i = 1, 2, \dots, m$. From (2.1) it follows that $0 \in \overline{co} \left(\bigcup_{i=1}^m \partial F(c, y_i) \right)$. Since $\|y_i - c\| = r > 0$ we have

$$\partial F(c, y_i) = \{x^* \in X^* : \|x^*\| = 1, \langle y_i - c, x^* \rangle = r\} = J(y_i - c).$$

As noted above J is a single-valued mapping, hence $J(y_i - c)$ consists of a unique point, say f_i . Therefore $0 \in \overline{co} \{f_1, f_2, \dots, f_m\} = co \{f_1, f_2, \dots, f_m\}$, and so there exist

non-negative numbers $\alpha_1, \alpha_2, \dots, \alpha_m$ such that $\sum_{i=1}^m \alpha_i = 1$ and $0 = \sum_{i=1}^m \alpha_i f_i$. Without loss of generality one can assume that all $\alpha_i, i = 1, 2, \dots, m$ are positive. It is a simple verification that these data also satisfy the conditions (i), (ii) above. The proof is complete.

Remark 2.2. We shall be interested mainly in the case $X = \ell_p$ ($p > 1$). For this purpose it is more convenient to use the following “scaled” version of the mapping J : for $x \in X$ and $p \in (1, \infty)$ we define $J(x) := \{x^* \in X^* : \langle x, x^* \rangle = \|x\| \cdot \|x^*\| = \|x\|^p\}$. Obviously for f_i, y_i and c as in Proposition 2.1, $\|y_i - c\|^{p-1} \cdot f_i \in J(y_i - c)$.

Remark 2.3. It is well-known that spaces ℓ_p ($1 < p < \infty$) are both uniformly convex and uniformly smooth. Hence the mapping J is single-valued. Also it is weakly sequentially continuous in the following sense: if $\{x_n\}$ converges weakly to x in ℓ_p , then $\{J(x_n)\}$ converges weakly to $J(x)$ in ℓ_q ($q = \frac{p}{p-1}$).

Remark 2.4. Clearly if $x = (x_1, x_2, \dots, x_n, \dots) \in \ell_p$ ($1 < p < \infty$), then $J(x) = (\text{sgn}(x_1) \cdot |x_1|^{p-1}, \text{sgn}(x_2) \cdot |x_2|^{p-1}, \dots, \text{sgn}(x_n) \cdot |x_n|^{p-1}, \dots)$. Hence in the situation of Proposition 2.1 with $X = \ell_p$ we have

$$f_i = \frac{J(y_i - c)}{\|y_i - c\|^{p-1}} = \frac{J(y_i - c)}{r^{p-1}}.$$

3. The results

For our use later it is convenient to formulate two auxiliary inequalities in the following lemma.

Lemma 3.1. *Let a and b be two real numbers. Then*

- (i) $\left| \text{sgn}(a) \cdot |a|^{p-1} - \text{sgn}(b) \cdot |b|^{p-1} \right| \leq 2^{2-p} \cdot |a - b|^{p-1}$, *provided* $1 < p \leq 2$;
- (ii) $|a|^p + |b|^p - p \cdot (a \cdot \text{sgn}(b) \cdot |b|^{p-1} + b \cdot \text{sgn}(a) \cdot |a|^{p-1}) \leq |a - b|^p$, *if* $p > 2$.

The proof is standard, and we shall omit it.

Theorem 3.2. *Let A be an extremal set in an ℓ_p space with $1 < p < \infty$. Then we have $\alpha(A) = d(A)$.*

Here $\alpha(A)$ denotes the Kuratowski measure of non-compactness of A .

Proof. We may assume $r_{\ell_p}(A) = 1$. Then for each integer number $n \geq 2$ we have $\bigcap_{x \in A} B(x, 1 - \frac{1}{n}) = \emptyset$, where $B(x, r)$ denotes the closed ball centered at x with radius r which

is weakly compact since ℓ_p is reflexive. Hence there exist $x_{t_{n-1}+1}, x_{t_{n-1}+2}, \dots, x_{t_n}$ in A such that $\bigcap_{i=t_{n-1}+1}^{t_n} B(x_i, 1 - \frac{1}{n}) = \emptyset$ (with convention $t_1 = 0$).

Setting $A_n := \{x_{t_{n-1}+1}, x_{t_{n-1}+2}, \dots, x_{t_n}\}$ we denote the Chebyshev center of A_n in ℓ_p by c_n and let $r_n := r_{\ell_p}(A_n)$, then $r_n > 1 - \frac{1}{n}$. In view of Proposition 2.1 one can find $y_{s_{n-1}+1}, y_{s_{n-1}+2}, \dots, y_{s_n}$ in A_n , continuous linear functionals $f_{s_{n-1}+1}, f_{s_{n-1}+2}, \dots, f_{s_n}$ on ℓ_p and positive numbers $\alpha_{s_{n-1}+1}, \alpha_{s_{n-1}+2}, \dots, \alpha_{s_n}$ (with convention $s_1 = 0$) such that:

- (i) $\langle y_i - c_n, f_i \rangle = r_n$ for $i = s_{n-1} + 1, s_{n-1} + 2, \dots, s_n$;
- (ii) $\|f_i\| = 1$, $i = s_{n-1} + 1, s_{n-1} + 2, \dots, s_n$;
- (iii) $\sum_{i=s_{n-1}+1}^{s_n} \alpha_i f_i = 0$, $\sum_{i=s_{n-1}+1}^{s_n} \alpha_i = 1$.

Setting $A_\infty := \{y_{s_{n-1}+1}, y_{s_{n-1}+2}, \dots, y_{s_n}\}_{n=2}^\infty$ we claim that $\alpha(A_\infty) = d(A)$. Suppose on the contrary $\alpha(A_\infty) < d(A)$. Then one can choose $\varepsilon_0 \in (0, d(A))$ satisfying $\alpha(A_\infty) \leq d(A) - \varepsilon_0$, and so subsets $D_1, D_2, \dots, D_m$ of ℓ_p with $d(D_i) \leq d(A) - \varepsilon_0$ for every $i = 1, 2, \dots, m$, such that $A_\infty \subset \bigcup_{i=1}^m D_i$. There exists at least one set among $D_1, D_2, \dots, D_m$, say D_1 with the property that there are infinitely many n satisfying

$$\sum_{i \in J_n} \alpha_i \geq \frac{1}{m} \quad (3.1)$$

where

$$J_n := \{i \in I_n := \{s_{n-1} + 1, s_{n-1} + 1, \dots, s_n\} : y_i \in D_1\}.$$

We shall estimate the sum $T_n := \sum_{i,j \in I_n} \alpha_i \alpha_j \langle y_i - y_j, f_i - f_j \rangle$. We have

$$\begin{aligned} T_n &= \sum_{i,j \in I_n} \alpha_i \alpha_j \langle (y_i - c_n) - (y_j - c_n), f_i - f_j \rangle = \\ &= \sum_{i,j \in I_n} \alpha_i \alpha_j \left[\langle y_i - c_n, f_i \rangle + \langle y_j - c_n, f_j \rangle - \langle y_i - c_n, f_j \rangle - \langle y_j - c_n, f_i \rangle \right] = \\ &= 2 \cdot \sum_{i,j \in I_n} \alpha_i \alpha_j r_n - \sum_{i \in I_n} \alpha_i \langle y_i - c_n, \sum_{j \in I_n} \alpha_j f_j \rangle - \sum_{j \in I_n} \alpha_j \langle y_j - c_n, \sum_{i \in I_n} \alpha_i f_i \rangle = \\ &= 2r_n - 2 \cdot \sum_{i \in I_n} \alpha_i \langle y_i - c_n, 0 \rangle = 2r_n \end{aligned} \quad (3.2)$$

As noted in Remark 2.4

$$f_i = \frac{J(y_i - c_n)}{\|y_i - c_n\|^{p-1}} = \frac{J(y_i - c_n)}{r_n^{p-1}}, \quad \forall i \in I_n.$$

Therefore for all $i, j \in I_n$

$$\langle y_i - y_j, f_i - f_j \rangle = \frac{1}{r_n^{p-1}} \cdot \langle (y_i - c_n) - (y_j - c_n), J(y_i - c_n) - J(y_j - c_n) \rangle.$$

1) *The case* $1 < p \leq 2$: Since the Jung constant $J(\ell_p) = \frac{1}{\sqrt[p]{2}}$, thus $d(A) = \sqrt[p]{2}$. Applying part (i) of Lemma 3.1 coordinatewise to the above expression and remembering $q = \frac{p}{p-1}$ one gets

$$\langle (y_i - c_n) - (y_j - c_n), J(y_i - c_n) - J(y_j - c_n) \rangle \leq 2^{2-p} \cdot \|(y_i - c_n) - (y_j - c_n)\|^p.$$

Hence

$$\begin{aligned} T_n &\leq \frac{1}{r_n^{p-1}} \sum_{i,j \in I_n} \alpha_i \cdot \alpha_j \cdot 2^{2-p} \cdot \|y_i - y_j\|^p = \\ &= \frac{2^{2-p}}{r_n^{p-1}} \cdot \left(\sum_{i,j \in J_n} \alpha_i \cdot \alpha_j \cdot \|y_i - y_j\|^p + \sum_{(i,j) \in I_n^2 \setminus J_n^2} \alpha_i \cdot \alpha_j \cdot \|y_i - y_j\|^p \right) \leq \\ &\leq \frac{2^{2-p}}{r_n^{p-1}} \cdot \left[(\sqrt[p]{2} - \varepsilon_0)^p \cdot \sum_{i,j \in J_n} \alpha_i \cdot \alpha_j + 2^{\frac{p}{q}} \cdot \left(1 - \sum_{i,j \in J_n} \alpha_i \cdot \alpha_j \right) \right] = \\ &= \frac{2^{2-p} \cdot 2^{\frac{p}{q}}}{r_n^{p-1}} - \frac{2^{2-p}}{r_n^{p-1}} \cdot \left(2^{\frac{p}{q}} - (\sqrt[p]{2} - \varepsilon_0)^p \right) \cdot \left(\sum_{i,j \in J_n} \alpha_i \cdot \alpha_j \right) \leq \\ &\text{(for all } n \text{ satisfying (3.1))} \\ &\leq \frac{2^{2-p} \cdot 2^{\frac{p}{q}}}{r_n^{p-1}} - \frac{2^{2-p}}{r_n^{p-1}} \cdot \left[2^{\frac{p}{q}} - (\sqrt[p]{2} - \varepsilon_0)^p \right] \cdot \frac{1}{m^2} = \\ &= \frac{2}{r_n^{p-1}} - \frac{2^{2-p}}{m^2 \cdot r_n^{p-1}} \cdot \left[2^{\frac{p}{q}} - (\sqrt[p]{2} - \varepsilon_0)^p \right] \end{aligned} \tag{3.3}$$

Comparing (3.2) and (3.3) we obtain

$$2 \cdot r_n^p \leq 2 - \frac{2^{2-p}}{m^2} \cdot \left[2^{\frac{p}{q}} - (\sqrt[p]{2} - \varepsilon_0)^p \right]$$

for all n satisfying (3.1). Since there infinitely many such n and $\lim_{n \rightarrow \infty} r_n = 1$, we come to a contradiction.

2) *The case $p > 2$:* Since the Jung constant $J(\ell_p) = \frac{1}{\sqrt[p]{2}}$, thus $d(A) = \sqrt[p]{2}$. Writting $y_i - c_n = (z_1^i, z_2^i, \dots)$, $J(y_i - c_n) = (v_1^i, v_2^i, \dots)$ with $v_j^i = \text{sgn}(z_j^i) \cdot \|z_j^i\|^{p-1}$, $j = 1, 2, \dots$ and applying part (ii) of Lemma 3.1 coordinatewise to $\langle (y_i - c_n) - (y_j - c_n), J(y_i - c_n) - J(y_j - c_n) \rangle$ we obtain

$$\begin{aligned} \langle (y_i - c_n) - (y_j - c_n), J(y_i - c_n) - J(y_j - c_n) \rangle &= \sum_{k=1}^{\infty} (z_k^i - z_k^j) \cdot (v_k^i - v_k^j) \leq \\ &\leq \sum_{k=1}^{\infty} \|z_k^i - z_k^j\|^p + (p-1) \cdot \sum_{k=1}^{\infty} (z_k^i \cdot v_k^j + z_k^j \cdot v_k^i), \end{aligned}$$

and therefore after summing up

$$\begin{aligned} r_n^{p-1} \cdot T_n &\leq \sum_{i,j \in I_n} \alpha_i \cdot \alpha_j \cdot \|y_i - y_j\|^p + (p-1) \cdot \sum_{i,j \in I_n} \alpha_i \cdot \alpha_j \cdot \sum_{k=1}^{\infty} (z_k^i \cdot v_k^j + z_k^j \cdot v_k^i) = \\ &= \sum_{i,j \in I_n} \alpha_i \cdot \alpha_j \cdot \|y_i - y_j\|^p + 2 \cdot (p-1) \cdot \sum_{k=1}^{\infty} \left(\sum_{i \in I_n} \alpha_i \cdot z_k^i \right) \cdot \left(\sum_{j \in I_n} \alpha_j \cdot v_k^j \right) = \\ &= \sum_{i,j \in I_n} \alpha_i \cdot \alpha_j \cdot \|y_i - y_j\|^p. \end{aligned}$$

The last equality follows from $\sum_{j \in I_n} \alpha_j \cdot v_k^j = 0$, $k = 1, 2, \dots$, which in turn can be deduced from the condition $\sum_{j \in I_n} \alpha_j \cdot f_j = 0$, or equivalently $\sum_{j \in I_n} \alpha_j \cdot J(y_j - c_n) = 0$. Thus in view of (3.2)

$$\begin{aligned} 2 \cdot r_n^p &\leq \sum_{i,j \in I_n} \alpha_i \cdot \alpha_j \cdot \|y_i - y_j\|^p = \\ &= \sum_{i,j \in J_n} \alpha_i \cdot \alpha_j \cdot \|y_i - y_j\|^p + \sum_{(i,j) \in I_n^2 \setminus J_n^2} \alpha_i \cdot \alpha_j \cdot \|y_i - y_j\|^p \leq \\ &\leq (\sqrt[p]{2} - \varepsilon_0)^p \cdot \left(\sum_{i,j \in J_n} \alpha_i \cdot \alpha_j \right) + 2 \cdot \left(1 - \sum_{i,j \in J_n} \alpha_i \cdot \alpha_j \right) = \end{aligned}$$

$$\begin{aligned}
&= 2 - \left[2 - (\sqrt[p]{2} - \varepsilon_0)^p \right] \cdot \left(\sum_{i,j \in J_n} \alpha_i \cdot \alpha_j \right) \leq \\
&\text{(for all } n \text{ satisfying (3.1))} \\
&\leq 2 - \left[2 - (\sqrt[p]{2} - \varepsilon_0)^p \right] \cdot \frac{1}{m^2},
\end{aligned}$$

a contradiction, because $\lim_{n \rightarrow \infty} r_n^p = 1$ and there are infinitely many n satisfying (3.1).

One concludes that $\alpha(A_\infty) = d(A)$, and hence $\alpha(A) = d(A)$.

The proof of Theorem 3.2 is complete.

As an immediate consequence one obtains an extension of Gulevich's result for ℓ_p spaces.

Corollary 3.3 (cf. [3]). *Let A be a relatively compact set in an ℓ_p space with $d(A) > 0$. Then*

$$\begin{aligned}
r_{\ell_p}(A) &< \frac{1}{\sqrt[p]{2}} \cdot d(A), & 1 < p \leq 2; \\
r_{\ell_p}(A) &< \frac{1}{\sqrt[p]{2}} \cdot d(A), & 2 < p < \infty.
\end{aligned}$$

Theorem 3.4. *Let A be an extremal set in a given ℓ_p space with $1 < p < \infty$. Then for every $\varepsilon \in (0, d(A))$, every positive integer m , there exists an m -simplex Δ with vertices in A such that each edge of Δ has length not less than $d(A) - \varepsilon$.*

Proof. We shall assume $r_{\ell_p}(A) = 1$. From the proof of Theorem 3.2 we derived a sequence $\{y_{s_{n-1}+1}, y_{s_{n-1}+2}, \dots, y_{s_n}\}_{n=2}^\infty$ in A , a sequence of continuous linear functionals $\{f_{s_{n-1}+1}, f_{s_{n-1}+2}, \dots, f_{s_n}\}_{n=2}^\infty$ in ℓ_q and a sequence of positive numbers $\{\alpha_{s_{n-1}+1}, \alpha_{s_{n-1}+2}, \dots, \alpha_{s_n}\}_{n=2}^\infty$ (with convention $s_1 = 0$) such that:

- (i) $\langle y_i - c_n, f_i \rangle = r_n$ for $i \in I_n := \{s_{n-1} + 1, s_{n-1} + 2, \dots, s_n\}$;
- (ii) $\|f_i\| = 1$, $i \in I_n$;
- (iii) $\sum_{i \in I_n} \alpha_i f_i = 0$, $\sum_{i \in I_n} \alpha_i = 1$,

where $c_n \in \ell_p$, and $r_n \in (1 - \frac{1}{n}, 1]$. Also we have

$$\begin{aligned}
2.r_n^p &\leq 2^{2-p} \cdot \sum_{i,j \in I_n} \alpha_i \cdot \alpha_j \cdot \|y_i - y_j\|^p, & 1 < p \leq 2; \\
2.r_n^p &\leq \sum_{i,j \in I_n} \alpha_i \cdot \alpha_j \cdot \|y_i - y_j\|^p, & 2 < p < \infty.
\end{aligned}$$

1) *The case* $1 < p \leq 2$. We denote by

$$\begin{aligned} T_{nj} &:= 2^{2-p} \cdot \sum_{i \in I_n} \alpha_i \cdot \|y_i - y_j\|^p, \\ S_n &:= \{j \in I_n : T_{nj} \geq 2 \cdot r_n^p \cdot (1 - \sqrt{1 - r_n^p})\}, \\ \lambda_n &:= \sum_{i \in I_n \setminus S_n} \alpha_i. \end{aligned}$$

We have

$$\begin{aligned} 2 \cdot r_n^p &\leq 2^{2-p} \cdot \sum_{i, j \in I_n} \alpha_i \cdot \alpha_j \cdot \|y_i - y_j\|^p = \\ &= \sum_{j \in S_n} \sum_{i \in I_n} 2^{2-p} \cdot \alpha_i \cdot \alpha_j \cdot \|y_i - y_j\|^p + \sum_{j \in I_n \setminus S_n} \sum_{i \in I_n} 2^{2-p} \cdot \alpha_i \cdot \alpha_j \cdot \|y_i - y_j\|^p \leq \\ &\leq 2^{2-p} \cdot 2^{\frac{p}{q}} \cdot \sum_{j \in S_n} \alpha_j + \sum_{j \in I_n \setminus S_n} \alpha_j \cdot 2 \cdot r_n^p \cdot (1 - \sqrt{1 - r_n^p}) = \\ &= 2 \cdot (1 - \lambda_n) + 2 \cdot \lambda_n \cdot r_n^p \cdot (1 - \sqrt{1 - r_n^p}) = \\ &= 2 - 2 \cdot \lambda_n \cdot (1 - r_n^p + r_n^p \cdot \sqrt{1 - r_n^p}) \leq \\ &\leq 2 - 2 \cdot \lambda_n \cdot \sqrt{1 - r_n^p}. \end{aligned}$$

Hence

$$\lambda_n \leq \sqrt{1 - r_n^p} \rightarrow 0$$

as $n \rightarrow \infty$. Thus

$$\lim_{n \rightarrow \infty} \left(\sum_{i \in S_n} \alpha_i \right) = \lim_{n \rightarrow \infty} (1 - \lambda_n) = 1.$$

On the other hand

$$\begin{aligned} \sum_{i, j \in I_n} \alpha_i \cdot \alpha_j \cdot \|y_i - y_j\|^p &= \sum_{i \neq j \in I_n} \alpha_i \cdot \alpha_j \cdot \|y_i - y_j\|^p \leq \\ &\leq 2^{\frac{p}{q}} \cdot \left(1 - \left(\sum_{i \in I_n} \alpha_i^2 \right) \right). \end{aligned}$$

So

$$2.r_n^p \leq 2^{2-p}.2^{\frac{p}{q}} \cdot \left[1 - \left(\sum_{i \in I_n} \alpha_i^2 \right) \right] \leq 2.(1 - \alpha_i^2)$$

for each $i \in I_n$. Therefore $\alpha_i \leq \sqrt{1 - r_n^p} \rightarrow 0$ as $n \rightarrow \infty$. One concludes that

$$|S_n| \longrightarrow \infty \quad \text{as} \quad n \rightarrow \infty.$$

This implies that for each positive integer m one can choose n sufficiently large so that $|S_n| > m$. For $j \in S_n$ put

$$\begin{aligned} S_n(y_j) &:= \left\{ i \in I_n : \|y_i - y_j\|^p \geq 2^{\frac{p}{q}} \cdot \left(1 - \frac{1}{\sqrt[4]{n}} \right) \right\}, \\ \hat{S}_n(y_j) &:= \{y_i : i \in S_n(y_j)\}. \end{aligned}$$

As $j \in S_n$ we have

$$\begin{aligned} 2.\left(1 - \frac{1}{n}\right)^p \cdot \left(1 - \sqrt{1 - \left(1 - \frac{1}{n}\right)^p}\right) &\leq 2.r_n^p \cdot \left(1 - \sqrt{1 - r_n^p}\right) \leq \\ &\leq 2^{2-p} \cdot \sum_{i \in I_n} \alpha_i \cdot \|y_i - y_j\|^p = \\ &= 2^{2-p} \cdot \sum_{i \in S_n(y_j)} \alpha_i \cdot \|y_i - y_j\|^p + 2^{2-p} \cdot \sum_{i \in I_n \setminus S_n(y_j)} \alpha_i \cdot \|y_i - y_j\|^p \leq \\ &\leq 2^{2-p} \cdot 2^{\frac{p}{q}} \cdot \left(1 - \sum_{i \in I_n \setminus S_n(y_j)} \alpha_i\right) + 2^{2-p} \cdot 2^{\frac{p}{q}} \cdot \left(1 - \frac{1}{\sqrt[4]{n}}\right) \cdot \left(\sum_{i \in I_n \setminus S_n(y_j)} \alpha_i\right) = \\ &= 2 - \frac{2}{\sqrt[4]{n}} \cdot \left(\sum_{i \in I_n \setminus S_n(y_j)} \alpha_i\right). \end{aligned}$$

Hence

$$\begin{aligned} \sum_{i \in I_n \setminus S_n(y_j)} \alpha_i &< \left[1 - \left(1 - \frac{1}{n}\right)^p \cdot \left(1 - \sqrt{1 - \left(1 - \frac{1}{n}\right)^p}\right) \right] \cdot \sqrt[4]{n} = \\ &= \left[1 - \left(1 - \frac{1}{n}\right)^p \right] \cdot \sqrt[4]{n} + \left(1 - \frac{1}{n}\right)^p \cdot \sqrt{1 - \left(1 - \frac{1}{n}\right)^p} \cdot \sqrt[4]{n} < \\ &< \frac{p}{\sqrt[4]{n^3}} + \sqrt{\frac{p}{\sqrt{n}}} < \\ &< \frac{2.p}{\sqrt[4]{n}} \end{aligned} \tag{3.4}$$

and so

$$\sum_{i \in S_n(y_j)} \alpha_i > 1 - \frac{2 \cdot p}{\sqrt[4]{n}} \quad (3.5)$$

Now for a given positive integer m we choose n sufficiently large such that

$$|S_n| > m \quad \text{and} \quad \frac{2 \cdot p \cdot m}{\sqrt[4]{n}} < 1.$$

We claim that for every choice of $i_1, i_2, \dots, i_m \in S_n$ we have

$$\bigcap_{k=1}^m S_n(y_{i_k}) \neq \emptyset \quad (3.6)$$

Indeed, otherwise $\bigcap_{k=1}^m S_n(y_{i_k}) = \emptyset$ would imply

$$S_n(y_{i_1}) \subset I_n \setminus \left(\bigcap_{k=2}^m S_n(y_{i_k}) \right) = \bigcup_{k=2}^m \left(I_n \setminus S_n(y_{i_k}) \right).$$

Consequently by (3.4) and (3.5)

$$\begin{aligned} 1 - \frac{2 \cdot p}{\sqrt[4]{n}} &< \sum_{\nu \in S_n(y_{i_1})} \alpha_\nu \leq \\ &\leq \sum_{k=2}^m \sum_{\nu \in I_n \setminus S_n(y_{i_k})} \alpha_\nu < (m-1) \cdot \frac{2 \cdot p}{\sqrt[4]{n}}, \end{aligned}$$

or $1 < \frac{2 \cdot p \cdot m}{\sqrt[4]{n}}$, a contradiction to the choice of n .

Furthermore from (3.6) it follows that if $1 \leq k \leq m$ and $i_1, i_2, \dots, i_k \in S_n$, then

$$\bigcap_{\nu=1}^k \hat{S}_n(y_{i_\nu}) \neq \emptyset.$$

With m and n as above let's fix some $j \in S_n$. Setting $z_1 := y_j$ we take consecutively $z_2 \in \hat{S}_n(z_1)$, $z_3 \in \hat{S}_n(z_1) \cap \hat{S}_n(z_2)$, $\dots$, $z_{m+1} \in \bigcap_{k=1}^m \hat{S}_n(z_k)$. Obviously

$$\|z_i - z_j\|^p \geq 2^{\frac{p}{q}} \cdot \left(1 - \frac{1}{\sqrt[4]{n}}\right)$$

for all $i \neq j$ in $\{1, 2, \dots, m+1\}$. Now for a given $\varepsilon \in (0, \sqrt[p]{2})$ one can choose n sufficiently large as above, and moreover so that

$$2^{\frac{p}{q}} \cdot \left(1 - \frac{1}{\sqrt[p]{n}}\right) \geq \left(\sqrt[p]{2} - \varepsilon\right)^p.$$

One obtains an m -simplex formed by $z_1, z_2, \dots, z_{m+1}$, whose edges have length not less than $\left(\sqrt[p]{2} - \varepsilon\right)$, as claimed.

2) *The case $2 < p < \infty$* can be proceeded in the same way just with replacing those T_{nj} and S_n in the first case suitably, *i.e.* as

$$T_{nj} := \sum_{i \in I_n} \alpha_i \cdot \|y_i - y_j\|^p,$$

$$S_n := \{j \in I_n : T_{nj} \geq 2 \cdot r_n^p \cdot (1 - \sqrt[p]{1 - r_n^p})\}.$$

The proof of Theorem 3.4 is complete.

References

- [1] Amir, D., *On Jung's constant and related constants in normed linear spaces*, Pacific J. Math., **118** (1985), 1–15.
- [2] Danzer, L., Grünbaum, B., Klee, V., *Helly's theorem and its relatives*, Proc. Symp. Pure Math., Amer. Math. Soc., **7** (1963), 101–180.
- [3] Gulevich, N. M., *The radius of a compact set in a Hilbert space*, Zapiski Nauchnykh Seminarov LOMI im. V. A. Steklova, AN SSSR, vol. **164** (1988), 157–158 (Russian).
- [4] Ioffe, A. D., Tikhomirov, V. M., *Theory of Extremal Problems*, Moscow: Nauka, 1974 (Russian).
- [5] Jung, H. W. E., *Über die kleinste Kugel, die eine räumliche Figur einschliesst*, J. Reine Angew. Math., **123** (1901), 241–257.
- [6] NguyenKhac, V., NguyenVan, K., *A characterization of extremal sets in Hilbert spaces*, arXiv: math. MG/0203190.
- [7] Pichugov, S. A., *Jung's constants of L_p space*, Math. Zametki, **43** (1988), 604–614 (Russian).

- [8] Routledge, N. A., *A result in Hilbert space*, Quart. J. Math., Oxford, (2) **3** (1952), N^o 9, 12–18.